

Publication list

Sahin H, Trautwein C, Wasmuth HE. Functional role of chemokines in liver disease models. *Nat Rev Gastroenterol Hepatol*. 2010;7:682-690.

Berres ML, Koenen RR, Rueland A, Zaldivar MM, Heinrichs D, **Sahin H**, Schmitz P, Streetz KL, Berg T, Gassler N, Weiskirchen R, Proudfoot A, Weber C, Trautwein C, Wasmuth HE. Antagonism of the chemokine Ccl5 ameliorates experimental liver fibrosis in mice. *J Clin Invest*. 2010;120:4129-4140.

Sahin H, Berres ML, Wasmuth HE. Therapeutic potential of chemokine receptor antagonists for liver disease. *Expert Rev Clin Pharmacol*. 2011;4:503-513.

Sahin H, Borkham-Kamphorst E, Kuppe C, Zaldivar MM, Grouls C, Al-samman M, Nellen A, Schmitz P, Heinrichs D, Berres ML, Doleschel D, Scholten D, Weiskirchen R, Moeller MJ, Kiessling F, Trautwein C, Wasmuth HE. Chemokine Cxcl9 attenuates liver fibrosis-associated angiogenesis in mice. *Hepatology*. 2012;55:1610-1619.

Sahin H, Trautwein C, Wasmuth HE. TLR4 stresses the liver. *Gut*. 2012;61:1241-1242.

Zaldivar MM, Berres ML, **Sahin H**, Nellen A, Heinrichs D, Schmitz P, Gassler N, Streetz KL, Trautwein C, Wasmuth HE. The chemokine receptor CXCR3 limits injury after acute toxic liver damage. *Lab Invest*. 2012;92:724-734.

Nellen A, Heinrichs D, Berres ML, **Sahin H**, Schmitz P, Proudfoot AE, Trautwein C, Wasmuth HE. Interference with oligomerization and glycosaminoglycan binding of the chemokine CCL5 improves experimental liver injury. *PLoS One*. 2012;7:e36614.

Sahin H, Tholema N, Petersen W, Raschke MJ, Stange R. Impaired biomechanical properties correlate with neoangiogenesis as well as VEGF and MMP-3 expression during rat patellar tendon healing. *J Orthop Res*. 2012;30:1952-1957.

Wiemuth D, **Sahin H**, Falkenburger BH, Lefevre CM, Wasmuth HE, Grunder S. BASIC--a bile acid-sensitive ion channel highly expressed in bile ducts. *FASEB J*. 2012;26:4122-4130.

Sahin H, Borkham-Kamphorst E, do ON, Berres ML, Kaldenbach M, Schmitz P, Weiskirchen R, Liedtke C, Streetz KL, Maedler K, Trautwein C, Wasmuth HE. Proapoptotic effects of the chemokine, CXCL 10 are mediated by the noncognate receptor TLR4 in hepatocytes. *Hepatology*. 2013;57(2):797-805.

Wiemuth D, **Sahin H**, Lefevre CM, Wasmuth HE, Grunder S. Strong activation of bile acid-sensitive ion channel (BASIC) by ursodeoxycholic acid. *Channels (Austin)*. 2012;7.

Sahin H, Wasmuth HE. Chemokines in tissue fibrosis. *Biochim Biophys Acta*. 2012

Meurer SK, Alsamman M, **Sahin H**, Wasmuth HE, Kisseleva T, Brenner DA, Trautwein C, Weiskirchen R, Scholten D. Overexpression of Endoglin modulates TGF- β 1-signalling pathways in a novel immortalized mouse hepatic stellate cell line. *PLoS One*. 2013;8:e56116.

Stock MK, Hammerich L, do O NT, Berres ML, Alsamman M, Heinrichs D, Nellen A, Trautwein C, Tacke F, Wasmuth HE, **Sahin H**. Met-CCL5 modifies monocyte subpopulations during liver fibrosis regression. *Int J Clin Exp Pathol*. 2013;6:678-685.

Heinrichs D, Berres ML, Nellen A, Fischer P, Scholten D, Trautwein C, Wasmuth HE, **Sahin H**. The Chemokine CCL3 Promotes Experimental Liver Fibrosis in Mice. PLoS One. 2013; 8:e66106.

Sahin H & Stange R, Wieskötter B, Persigehl T, Ring J, Bremer C, Raschke MJ, Vieth V. In vivo monitoring of angiogenesis during tendon repair: a novel MRI-based technique in a rat patellar tendon model. Knee Surg Sports Traumatol Arthrosc. Knee Surg Sports Traumatol Arthrosc 2014.

Zimmermann HW, Sterzer V, **Sahin H**. CCR1 and CCR2 Antagonists. Curr Top Med Chem 2014;14:1539-1552.